

PATIENT DETAILS			
NAME		DATE OF BIRTH (DD/MM/YYYY)	
PHONE		EMAIL	
ADDRESS		HEALTH CARD NUMBER	
EMERGENCY CONTACT NAME		EMERGENCY CONTACT NUMBER	

CLINICAL DETAILS			
DIAGNOSIS		HEMOGLOBIN	g/l
WEIGHT (KG)		FERRITIN	ng/mL
Is patient pregnant, breastfeeding, or under the age of 18?		<input type="checkbox"/> No <input type="checkbox"/> Yes → Please prescribe Venofer instead as Monoferric is not currently approved for use in pregnancy/lactation or patients under age 18 in Canada. Please note that Venofer should not be given to pregnant women in the first trimester.	
Has patient received IV iron previously?		<input type="checkbox"/> No <input type="checkbox"/> Yes → Indicate if any reaction:	

PRESCRIPTION																			
<input type="checkbox"/> MONOFERRIC Simplified Monoferric Weight-Based Table <table border="1"> <thead> <tr> <th>Hb (g/L)</th> <th><50kg</th> <th>50-70kg</th> <th>≥70kg</th> </tr> </thead> <tbody> <tr> <td>≥100</td> <td>500mg</td> <td>1000mg</td> <td>1500mg</td> </tr> <tr> <td><100</td> <td>500mg</td> <td>1500mg</td> <td>2000mg</td> </tr> </tbody> </table> Doses that exceed the weight-based chart above, 20mg iron/kg body weight, or 1500mg, must be split into multiple doses separated by at least 7 days (Induction Dose). If the dose is not clearly specified, the product monograph administration guidelines will be followed.		Hb (g/L)	<50kg	50-70kg	≥70kg	≥100	500mg	1000mg	1500mg	<100	500mg	1500mg	2000mg	<input type="checkbox"/> VENOFER Simplified Venofer Dosing Table Max Dose for Treatment Regime = 1000mg Max Daily Dose = 300mg					
Hb (g/L)	<50kg	50-70kg	≥70kg																
≥100	500mg	1000mg	1500mg																
<100	500mg	1500mg	2000mg																
<input type="checkbox"/> FERINJECT Determine total iron requirements according to the table below. A single administration should not exceed 15 mg/kg/dose (max. 1000 mg). Base on total iron requirements, if an additional dose is required, do not administer before at least 7 days.																			
DOSE	DOSING REGIMEN	Weight	Hemoglobin																
<input type="checkbox"/> 500mg <input type="checkbox"/> 1000mg <input type="checkbox"/> 1500mg <input type="checkbox"/> 2000mg (induction) Total Number of Doses: _____ Interval: <input type="checkbox"/> 2 months <input type="checkbox"/> 3 months <input type="checkbox"/> 6 months <input type="checkbox"/> Other: _____	<input type="checkbox"/> 200mg IV every ____ week(s) for ____ doses <input type="checkbox"/> 300mg IV every ____ week(s) for ____ doses <input type="checkbox"/> Other: ____ mg IV every ____ week(s) for ____ doses	<table border="1"> <thead> <tr> <th></th> <th>< 10 g/dL</th> <th>10 to < 14 g/dL</th> <th>≥14 g/dL</th> </tr> </thead> <tbody> <tr> <td>< 35 kg</td> <td>500 mg IV</td> <td>500 mg IV</td> <td>500 mg IV</td> </tr> <tr> <td>35 kg - < 70 kg</td> <td>1500 mg IV</td> <td>1000 mg IV</td> <td>500 mg IV</td> </tr> <tr> <td>≥70 kg</td> <td>2000 mg IV</td> <td>1500 mg IV</td> <td>500 mg IV</td> </tr> </tbody> </table>		< 10 g/dL	10 to < 14 g/dL	≥14 g/dL	< 35 kg	500 mg IV	500 mg IV	500 mg IV	35 kg - < 70 kg	1500 mg IV	1000 mg IV	500 mg IV	≥70 kg	2000 mg IV	1500 mg IV	500 mg IV	
	< 10 g/dL	10 to < 14 g/dL	≥14 g/dL																
< 35 kg	500 mg IV	500 mg IV	500 mg IV																
35 kg - < 70 kg	1500 mg IV	1000 mg IV	500 mg IV																
≥70 kg	2000 mg IV	1500 mg IV	500 mg IV																

OTHER MEDICATIONS		
If the patient has a HISTORY of reaction to any Iron products, give the following medication IMMEDIATELY prior to the infusion: <input type="checkbox"/> Methylprednisolone 125mg IV x1 <input type="checkbox"/> Diphenhydramine 25-50 mg PO/IV <input type="checkbox"/> Acetaminophen 650 mg PO <input type="checkbox"/> Other:	<input type="checkbox"/> Our clinics follow a standardized protocol to manage reactions during our post-infusion. Please tick this box to indicate that you agree with the following protocol. If the patient has adverse reaction DURING/POST infusion, give: <input type="checkbox"/> Hydrocortisone 100mg IV <input type="checkbox"/> Methylprednisolone 125mg IV <input type="checkbox"/> Diphenhydramine 25-50mg PO/IV <input type="checkbox"/> Acetaminophen 650mg PO <input type="checkbox"/> Dimenhydrinate Gravol® 25-50mg PO/IV	Current infusion reaction protocol includes the use of these medications according to nurse's assessment.

ADDRESS		PHONE		FAX	
PRESCRIBER NAME		LICENSE NUMBER			
PRESCRIBER SIGNATURE		DATE (DD/MM/YYYY)			

All medications, IV solutions, and related supplies used in IV therapy are dispensed and provided by CareMed pharmacy in full compliance with applicable regulatory and safety standards.

To ensure product safety and integrity, CareMed Wellness Clinic does not accept or administer any medications or IV products sourced or filled by a pharmacy other than CareMed Pharmacy. All medications used in treatment must be dispensed directly through CareMed Pharmacy.